

The impact on cosmology of a primordial scaling field

Pedro G. Ferreira^{a*}

^a*Center for Particle Astrophysics, 301 Leconte Hall,
University of California, Berkeley, CA 94720*

Abstract. A scalar field with an exponential potential has the particular property that it is attracted into a solution in which its energy scales as the dominant component (radiation or matter) of the Universe, contributing a fixed fraction of the total energy density. We briefly discuss the dynamics of such a scalar field and its impact on Big Bang nucleosynthesis, the growth of large scale structure and abundance of damped Lyman- α systems at high redshift. Given the simplicity of the model, its theoretical motivation, and its success in matching observations, we argue that it should be taken on par with other currently viable models of structure formation.

The increasing quality of data from large scale surveys, cosmic microwave background experiments, high redshift observations, and other cosmological measures, has begun to seriously constrain the range of allowed models of structure formation. In particular the simplest models have been shown incapable of predicting the right level of fluctuations consistently on Gpc and Mpc scales. The alternatives invoke some form of fine tuning. This can be in the form of very special initial conditions so that the energy density in a cosmological constant, curvature, or some other exotic form of matter comes to dominate at late times, or it can be in the form of a non-clustering component that is motivated by standard particle physics but whose mass scale is difficult to reconcile with the known particle bestiary.

Given the the unsatisfactory collection of models one has to work with, it makes sense to find alternatives that might point us in the right direction for finding a compelling and observationally viable model of structure formation. It turns out that a general class of theories in particles physics (Kaluza Klein, Supergravity, or Superstring based) predict the existence of a weakly coupled scalar field, Φ , with a potential of the form

$$V(\Phi) = M_P^4 \exp\left(-\lambda \frac{\Phi}{M_P}\right), \quad \lambda \sim \mathcal{O}(1) \quad (1)$$

where M_P is the Planck mass, which has an attractor solution (and therefore is insensitive to initial conditions) and leads to a consistent level of fluctuations on all scales [1].

As shown in [2, 3, 4] a scalar field with an exponential potential (and $\lambda > 2$) will evolve in

*This work was done in collaboration with Michael Joyce. Support came from NSF Cooperative Agreement No. AST 9120005, PRAXIS XXI (Portugal) and CNRS (France)

such a way that its fractional energy density, Ω_Φ , is constant. In particular one can show that

$$\Omega_\Phi = \frac{n}{\lambda^2} \quad (2)$$

where n is 3 (4) in the matter (radiation) era. The competition between the friction due to the expansion of the universe and the inertia from the field rolling down the potential make it an attractor: if the energy density in Φ , ρ_Φ , is small compared to the total energy density, it will remain constant until it becomes relevant. On the other hand if ρ_Φ dominates, then it will evolve as $\rho_\Phi \propto a^{-\lambda^2}$ until it catches up with the background energy density. In the spirit of simplicity we shall assume that the scalar field has been in the attractor from very early times. In particular it contributed a non-negligible fraction to the relativistic background at the time of nucleosynthesis. This supplies us with a constraint on Ω_Φ today:

$$\Omega_\Phi < 0.15 - 0.2 \quad (3)$$

The fact that Ω_Φ is so small might lead one to think that such a scalar field has little or no impact on structure formation. However the fact that such matter makes up a constant fraction of the total energy density during the whole matter-dominated era make its effects considerable. Indeed one can show that for an $H_0 = 50(65) \text{ km s}^{-1} \text{ Mpc}^{-1}$, an $\Omega_\Phi = 0.08(0.12)$ suffices to reconcile the cosmic microwave background data on horizon size scales with the large scale structure data (see Figures 1 and 2). We shall call such a class of models Φ CDM.

The evolution of perturbations in the presence of the scalar field is simple to understand. On superhorizon scales there is the usual growing mode with $\delta_c \propto \tau^2$ (where δ_c is the density contrast in the CDM) and τ is conformal time. This is to be expected; the superhorizon evolution is insensitive to the “chemistry” of the matter and totally dominated by gravity. On sub-horizon scales in the radiation era, the Meszaros effect comes into play giving $\delta_c \propto \ln \tau$. The specific effect of the scalar field appears on subhorizon scales in the matter era. The perturbation in the scalar field itself has the approximate solution $\delta_\Phi \propto (1/\tau^{3/2})J_{3/2}(k\tau)$ (where J_ν is a Bessel function) which, when fed back into the equation for δ_c , gives an altered solution for the usual growing mode $\delta_c \propto \tau^{2-\epsilon}$ where

$$\epsilon = \frac{5}{2} \left(1 - \sqrt{1 - \frac{24}{25}\Omega_\Phi} \right) \quad (4)$$

This solution shows explicitly how even a small contribution from the scalar field can give a significant effect, as it acts all the way through the matter era. The expected suppression of $|\delta_c|^2$ for modes larger than k_{eq} is of order $(1 + z_{eq})^{-\epsilon}$, where k_{eq} is the wavenumber of the horizon size at radiation-matter equality. This last effect is reminiscent of the evolution of perturbations in a mixed dark matter (MDM) universe where one has component of matter, ρ_ν , which is collisionless for a period of time during the matter era.

There is, however a crucial difference between our model and MDM: the period of time during which perturbations are suppressed is shorter in MDM compared to Φ CDM. In both cases there is a wavenumber, k_{su} , which separates growing modes from damped modes. For Φ CDM, this scale is roughly the horizon, i.e., $k_{su} \propto 1/\tau$, while for MDM it is the free streaming scale, i.e., $k_{su} = 8a^{1/2}(m_\nu/10 \text{ eV})h \text{ Mpc}^{-1} \propto \tau$. Clearly in the latter case any given mode of δ_c will




Figure 1. Mass variance per unit $\ln k$ computed from the Boltzman code for different models compared with that inferred from a compilation of galaxy surveys.




Figure 2. Comparison of different model predictions to current experimental data. All models were COBE normalized and are labeled as in Figure 1.

eventually start to grow. In particular, modes around k_{eq} will already have started to undergo collapse.

Finally, we find Φ CDM has an interesting advantage over MDM. MDM models predict too little structure compared to that inferred from the Lyman- α measurements. From Figure 1 we can see that Φ CDM should fare better than MDM on very small scales. This is easy to understand: We argued that effectiveness of Φ CDM was mainly due to the fact that the scalar field free-streaming scale grows with time, while the massive neutrino free-streaming scale decays with time. We then need a larger amount of massive neutrinos to fit both COBE and the cluster abundances in the MDM model than the amount of scalar field in Φ CDM. On much smaller scales (the scales probed by Lyman- α systems), i.e., scales smaller than the massive neutrino free-streaming scale, perturbations in MDM should be more suppressed than in Φ CDM. This means Φ CDM should fare better than MDM with regards to the Lyman- α constraints.

References

- [1] P.G. Ferreira and M. Joyce, Phys. Rev. Lett. 79 (1997) 4740.
- [2] B. Ratra and J. Peebles, Phys. Rev. D 37 (1988) 3406.
- [3] E. Copeland, A. Liddle, and D. Wands, Ann. N. Y. Acad. Sci. 688 (1993) 647.
- [4] C. Wetterich, Astron. Astrophys. 301 (1995) 321.